

The extraction of g_A from finite volume systems: The long and short of it

Thomas D. Cohen

Department of Physics, University of Maryland, College Park 20742

UMPP#02-025

DOE/ER/40762-247

Due to a pion-pole contribution, the nucleon axial current matrix elements can be visualized in position space as becoming delocalized as the chiral limit is approached, with one-third of the current at distance of order $1/m_\pi$. However, this delocalization effect will not cause calculations of g_A in a finite system with standard boundary conditions (periodic for boson fields, antiperiodic for fermions) to have large finite volume effects. This is seen by calculating g_A using a nonstandard current with axial quantum numbers. The matrix elements of this new current lacks a pion pole and is not delocalized; however, its zero-momentum matrix element is identical to that of the standard axial current both for infinite and finite volumes.

The coupling of the nucleon to axial currents at zero momentum transfer is fixed by the coupling constant g_A . Due to chiral symmetry and the key role played by pions in the low-lying dynamics of the nucleon, understanding how g_A emerges from the underlying dynamics of QCD is essential if one wishes to understand how the low energy dynamics of the nucleon is connected to QCD. Attempts to calculate g_A in lattice QCD have given values which are well below the experimental value [1]. It has recently been argued that this failure may be due to very large finite volume effects [2]. One may ask whether there is a basis for understanding the emergence of large finite volume effects from the structure of QCD. This issue was brought into sharp focus in a recent paper by Jaffe [3]. Jaffe made an elegant argument on very general grounds that the spatial distribution of the axial current in the nucleon is necessarily spread over large distance as the chiral limit is approached, a phenomenon he denotes by “delocalization”. In particular he showed that as the quark mass goes to zero, one-third of the axial current is pushed off to infinite distance. For finite quark mass, this implies that of order one-third of the axial current can be found at a distance of order m_π^{-1} . It was suggested in Ref. [3] that this effect could lead to large finite volume effects. Intuitively it is not implausible that this delocalization of the axial current could lead to large finite volume effects in calculations of g_A : if the pion is light enough so that large amounts of the axial current are pushed to regions which are beyond the lattice volume one might expect that the lattice calculation will miss these contributions leading to substantial errors.

The purpose of this paper is to observe that although ref [3] is correct that one-third of the axial current is delocalized in the chiral limit, the intuition that this implies large finite volume effects is not (provided the standard boundary conditions of periodicity for bosons and anti-periodicity for fermions are imposed). To make this statement a bit more precise it is necessary to look a bit more carefully at the heuristic argument why one might expect large finite volume effects. The basic issue is the play off between the infinite volume and chiral limits. The most serious worry is that as the chiral limit

is approached, one-third of the axial current will be lost outside the box for any finite sized box. Thus to get convergence of g_A one needs to choose a box size significantly larger than one with a length scale set by $1/m_\pi$. Mathematically this would be reflected by nonuniform limits of the calculated value of g_A as one approaches the infinite spatial volume and chiral limits. Define g_A^{Vol} to be the matrix element of the axial current between nucleon states as calculated with fixed quark mass (which fixes m_π) in a finite spatial volume. The principal concern is that g_A^{Vol} calculated in the limit $V \rightarrow \infty, m_\pi \rightarrow 0$ will only be two thirds of g_A^{Vol} calculated in the limit $m_\pi \rightarrow 0, V \rightarrow \infty$. However as will be shown below, if the standard boundary conditions are imposed the limit should be uniform:

$$\lim_{V \rightarrow \infty, m_\pi \rightarrow 0} g_A^{\text{Vol}} = \lim_{m_\pi \rightarrow 0, V \rightarrow \infty} g_A^{\text{Vol}}. \quad (1)$$

Thus, the major intuitive concern about large finite volume effects can be put to rest. There is a secondary worry, however, that if much of the strength of the underlying integrals come from regions comparable to the box size, the dynamical effects induced by the boundary could alter this part of the distribution even if the strength is not actually lost. This could lead to finite volume effects which are larger than those for typical baryon matrix elements. This secondary concern has not been stated in precise form and, hence, it is hard to precisely contradict. However, it will also be shown below there is no reason based on the delocalization of current discussed in Ref. [3] to expect that finite volume effects to be larger for g_A than those for typical matrix elements.

Let us begin with a brief discussion of the demonstration that the axial current is delocalized. In a certain sense, the argument for the delocalization is quite old; the result is well known in the context of mean field chiral soliton and bag models of the nucleon [4]. In these models the nucleon is treated as a localized source and the pion field is treated classically. The localization of the source and the mean field treatments are typically justified from large N_c QCD arguments. The axial coupling constant is calculated by integrating the axial current over the nucleon and then matching to the general

Lorentz covariant expression for axial matrix elements assuming that the nucleon source is localized. The result is that as $m_\pi \rightarrow 0$ the integrals which give g_A receives one-third of its contribution from distances of order m_π , and as one approaches the chiral limit one-third of the integrated strength is pushed off to infinity. In a sense, the crux of the recent argument of Ref. [3] is that these well-known results in the context of mean field chiral models can be translated into a general model-independent result with essentially no change in structure.

To begin, consider the general structure of axial matrix elements of the nucleon in terms of form factors:

$$\begin{aligned} \langle N(\vec{p}', s', I'_3) | A_a^\mu(0) | N(\vec{p}, s, I_3) \rangle = \\ \bar{u}(\vec{p}', s', I'_3) \frac{\tau_a}{2} (g_A(q^2) \gamma^\mu \gamma_5 + h_A(q^2) q^\mu \gamma_5) u(\vec{p}, s, I_3), \end{aligned} \quad (2)$$

where $u(\vec{p}, s, I_3)$ is a free isospin one-half Dirac spinor with a third component of spin, s , and a third component of isospin, I_3 [5]. The form factors, $g_A(q^2)$ and $h_A(q^2)$, are generally referred to as axial and induced pseudoscalar form factors, respectively. The axial coupling constant, g_A , is defined as the axial form factor at zero momentum transfer. The key point is that the axial current acting on the vacuum can create a pion which implies that the induced pseudoscalar form factor has a pion pole contribution:

$$\begin{aligned} h_A(q^2) &= h_A^{pole}(q^2) + h_A^{short}(q^2) \\ h_A^{pole}(q^2) &= -2f_\pi g_{\pi NN}, \end{aligned} \quad (3)$$

where f_π is the pion decay constant (with the convention that $f_\pi \approx 93\text{MeV}$), $g_{\pi NN}$ is the pion-nucleon coupling constant, and $h_A^{short}(q^2)$ the non-pion-pole part of the induced pseudoscalar form factor. This non-pion-pole contribution to h_A is denoted as “short” to indicate the scale over which it varies substantially is the hadronic scale, $\Lambda \sim 1\text{GeV}$, which is much larger than m_π and hence corresponds to a spatial extent which is much smaller than $1/m_\pi$. The pion-pole contribution to the induced pseudoscalar coupling, however, varies substantially if q^2 changes on a scale comparable to m_π which implies contributions on a spatial extent of order $1/m_\pi$. This rapid variation of the pion pole term is the origin of the effective delocalization of the axial current.

Rather than recapitulate the demonstration of Ref. [3] in detail, an equivalent and somewhat simpler derivation of the same result will be presented here. Consider a spatial component of the axial current between spinors. (A spatial component is used since the temporal component vanishes at zero momentum transfer). At finite momentum transfer the results are frame dependent. It is natural to work in the Breit frame of ($\vec{p}_{initial} = -\vec{p}_{final}$). The Breit frame has the virtue of having momentum transfer but no energy transfer. Thus the Fourier transform of the matrix element in the Breit frame may be directly interpreted in terms of position space; the relative time

variable plays no role. Introducing the symbol $G(\vec{q})$ as the matrix element of the third spin and isospin components of the axial current between spin-up protons in the Breit frame and a three-momentum transfer of $\vec{Q}$, and using Eq. (3) and standard properties of Dirac spinors one has:

$$\begin{aligned} G(Q^2) &\equiv \langle N(\vec{Q}, 1/2, 1/2) | A_3^0(0) | N(-\vec{Q}, 1/2, 1/2) \rangle = \\ &\left(g_A(-|\vec{Q}|^2) \left(1 + \frac{|\vec{Q}|^2 - 2Q_3^2}{16M^2} \right) + h_A(-|\vec{Q}|^2) \frac{Q_3^2}{2M} \right) \\ &\quad \times \frac{\sqrt{Q^2 + 4M^2} + 2M}{4M}, \end{aligned} \quad (4)$$

where M is the nucleon mass. The choice of the third spin and isospin components and the proton spin up states was arbitrary and done for the purposes of specificity; other choices of components and spin and isospin projections of the state lead to identical results.

From Eq. (4) it is immediately apparent that $\lim_{\vec{Q} \rightarrow 0} G(\vec{Q}) = g_A$, providing that there is no singularity in the induced pseudoscalar form factor at $q^2 = 0$. For any finite m_π this is true and the limit both exists and is uniform in the sense that one can take the $\vec{Q} \rightarrow 0$ limit from any direction and will get the same answer. Because the limit $\vec{Q} \rightarrow 0$ limit is uniform, if we define an angular average of $G(\vec{Q})$,

$$\overline{G}(Q^2) \equiv \frac{1}{4\pi} \int d\Omega G(\vec{Q}), \quad (5)$$

then it is immediately clear that for any nonzero m_π ,

$$\lim_{Q \rightarrow 0} \overline{G}(Q^2) = g_A. \quad (6)$$

We wish to study $\overline{G}(Q^2)$ for both q and m_π well below the typical hadronic scale of $\Lambda \sim 1\text{GeV}$. Assuming $g_A(q^2)$ and $h_A(q^2)$ both vary over scales of order Λ , one sees from Eqs. (2), (3), (4) and (5) that

$$\overline{G}(Q^2) = g_A - \frac{f_\pi g_{\pi NN}}{3M} \frac{Q^2}{Q^2 + m_\pi^2} + \mathcal{O}(Q^2/\Lambda^2, m_\pi^2/\Lambda^2). \quad (7)$$

Imposing conservation of axial charge in the chiral limit on Eq. (3) yields the Goldberger-Treiman relation $g_A M = f_\pi g_{\pi NN} + \mathcal{O}(m_\pi^2/\Lambda^2)$ [6] so that

$$\overline{G}(q^2) = g_A \left(1 - \frac{1}{3} \frac{q^2}{q^2 + m_\pi^2} \right) + \mathcal{O}(q^2/\Lambda^2, m_\pi^2/\Lambda^2). \quad (8)$$

Note that the preceding equation is the leading order term in chiral expansion; while there are correction terms of order Q^2/Λ^2 and m_π^2/Λ^2 , Q^2/m_π^2 has been included to all orders.

The delocalization of axial charge discussed in Ref. [3] is clear from Eq. (8). The term proportional to $Q^2/(Q^2 +$

m_π^2) clearly contributes $1/3$ of the $Q = 0$ result. It is also clear that this term varies rapidly over momentum scales corresponding to spatial variations over long distance scales $\sim 1/m_\pi$. As $m_\pi \rightarrow 0$ the spatial distance over which this term contributes becomes arbitrarily large, *i.e.* the current becomes delocalized. To understand this as a spatial delocalization of the current in the chiral limit, one needs a method to visualize the current in position space. The natural way is to define a spatial axial current density $\rho_A(r)$ as the Fourier transform of $\overline{G}(q^2)$. If the current is delocalized in the chiral limit, then all of the positive moments of $\rho_A(r)$ will diverge. These moments are given by

$$\begin{aligned} \langle r^{2n} \rangle_A &= \frac{1}{4\pi} \int d^3r r^{2n} \rho_A(r) \\ &= \prod_{j=1,n} (2+4j) (-1)^n \frac{d^n \overline{G}(Q^2)}{d(Q^2)^n}. \end{aligned} \quad (9)$$

Using Eq. (8) one has

$$\begin{aligned} \langle r^{2n} \rangle_A &= \frac{g_A n! \prod_{j=1,n} (2+4j)}{3m_\pi^{2n}} (1 + \mathcal{O}(m_\pi^2/\lambda^2)) \\ &\text{for } n \geq 1. \end{aligned} \quad (10)$$

all of which diverges $\sim m_\pi^{-2n}$ as the limit $m_\pi \rightarrow 0$ is approached. This indicates a delocalized current in the chiral limit.

Thus, one-third of the axial current density can be visualized as being delocalized as the chiral limit is approached. While this represents an important insight into the physics, it is important to recall that the underlying dynamics depends on nothing beyond the fact that the axial current matrix element has a pion pole contribution. In this paper, the central issue is the relationship between the delocalization of the axial current and effects due to finite spatial volumes. Of course, real lattice calculations have a number of sources of errors including statistical and discretization errors as well as errors due to finite spatial and temporal extent. Here, the focus is on finite spatial volume effects. To isolate these effects it is useful to consider the behavior of continuum QCD on a finite three-dimensional volume with infinite temporal extent. From the prospect of lattice QCD this can be thought of as working with a sufficiently fine mesh, a sufficiently large number of lattice points in the time direction and a sufficiently large number of configurations so that discretization errors—errors associated with finite temporal extent—and statistical errors are all negligible.

The calculation of g_A in finite spatial volume QCD is straightforward in principle. We consider the three-dimensional space to be a cube of length L , and impose the boundary conditions that fermion fields are anti-periodic and boson fields are periodic. Since the time direction is infinite, one can define time evolution as being generated by a hermitian hamiltonian operator

whose eigenstates have a well-defined energy. These energy eigenstates are most easily treated as being box-normalized. Because these states are associated with fields which have well-defined properties under translation in the x , y , or z direction by L , the energy states are all eigenstates of the momentum operator with $\vec{p} = 2\pi L^{-1}(n_x \hat{x} + n_y \hat{y} + n_z \hat{z})$ (integer n 's) for bosons and with $\vec{p} = \pi L^{-1}((2n_x + 1)\hat{x} + (2n_y + 1)\hat{y} + (2n_z + 1)\hat{z})$ (integer n 's) for fermions. The box breaks rotational symmetry and hence angular momentum is no longer a good quantum number. However, the spin degree of freedom is still well defined for fermions in the sense that for every momentum there are two degenerate eigenstates. Thus the nucleon states can be labeled as $|\vec{p}, I_3, s\rangle$ with the normalization $\langle \vec{p}', I'_3, s' | \vec{p}, I_3, s \rangle = \delta_{I'_3, I_3} \delta_{s', s} \delta_{n'_x, n_x} \delta_{n'_y, n_y} \delta_{n'_z, n_z}$. One can define the finite volume version of g_A as the diagonal matrix element of the third space and isospace components of axial current between spin up proton states on minimum momentum appropriately normalized to the volume of the box:

$$\begin{aligned} g_A^{\text{Vol}} &= L^3 \langle \vec{p}, I_3, s | A_3^3(\vec{x}, t) | \vec{p}, I_3, s \rangle \\ &= \langle \vec{p}, I_3, s | \int_{\text{box}} d^3x A_3^3(\vec{x}, t) | \vec{p}, I_3, s \rangle, \end{aligned} \quad (11)$$

where the second form of Eq. (11) stems from translational invariance of the matrix element with these boundary conditions.

Before proceeding it is useful to distinguish between the method of calculating g_A for the finite spatial system discussed above as an integrated quantity in a state of good energy from the method actually used in lattice calculations. In lattice calculations one never calculates exact states of good energy. Nucleon matrix elements such as g_A are calculated as the ratio of a three-point correlation function to a two-point function with the times of the three currents well separated in time. As the separation in time between these currents becomes large, the system becomes dominated exponentially by the lowest energy state with quantum numbers which overlap those of the initial and final currents. As the time separation between all three currents goes to infinity the ratio of the three-point function to the two-point function asymptotically approaches the matrix element in the nucleon state considered above. Thus the matrix derived using the states of good energy are equivalent to the ratio of the three-point functions to the two-point functions used in lattice calculations proven that the times are arbitrarily long. If the times are not arbitrarily long there are finite corrections.

The issue of concern is whether the delocalization of the axial current for the infinite volume theory leads to large finite volume errors. It is easy to show that this is not the case by considering the matrix element of a new axial current operator. Suppose that we can find a current operator which in the infinite volume limit does

not suffer from the delocalization effect seen in the conventional axial current. If it can be shown that the integrated matrix element of this new current in the finite volume theory precisely equals g_A^{Vol} , then one can replace the original finite volume calculation of g_A with one based on the new current. Doing this has the virtue that delocalization does not occur in the new calculation and hence cannot be responsible for large finite volume effects. Since g_A^{Vol} equals the finite volume QCD matrix element of the new current integrated over space, it follows that g_A^{Vol} also cannot have a large finite volume effect due to delocalization. It is straightforward to find a current that has these properties. Since the delocalization is traceable to the pion pole, we need to find a current whose nucleon matrix elements lack such a pole.

Consider the current $A_a'^\mu$ defined by

$$A_a'^\mu \equiv \left(1 + \frac{\partial^\alpha \partial_\alpha}{m_\pi^2}\right) A_a^\mu, \quad (12)$$

where $\partial^\alpha \partial_\alpha$ is the D'Alembertian operator. Although this current has no well-defined chiral limit, it is well defined for any finite value of the quark mass and this is sufficient for the present purpose. The matrix elements of this current can easily be expressed in terms of the old one:

$$\begin{aligned} N(\vec{p}', s', I_3') |A_a'^\mu(0) |N(\vec{p}s, I_3)\rangle &= \\ \bar{u}(\vec{p}', s', I_3') \frac{1}{2} \langle \tau_a (g'_A(q^2) \gamma^\mu \gamma_5 + h'_A(q^2) q^\mu \gamma_5) u(\vec{p}, s, I_3) & \\ g'_A(q^2) = \left(1 - \frac{q^2}{m_\pi^2}\right) g_A(q^2) & \\ h'_A(q^2) = \left(1 - \frac{q^2}{m_\pi^2}\right) h_A(q^2). & \end{aligned} \quad (13)$$

It is clear from this form that as $q \rightarrow 0$, $g'_A \rightarrow g_A$ and $h'_A \rightarrow 0$. For the infinite volume case, one can define the function $\overline{G}'(Q^2)$ in direct analogy to $\overline{G}(Q^2)$,

$$\begin{aligned} \overline{G}'(Q^2) &\equiv \frac{1}{4\pi} \int d\Omega G'(\vec{Q}) \\ G'(\vec{Q}) &\equiv \langle N(\vec{Q}, 1/2, 1/2) | A_3'^3(0) | N(-\vec{Q}, 1/2, 1/2) \rangle. \end{aligned} \quad (14)$$

Inserting the matrix elements for A' from Eq. (13) into the definition for $\overline{G}'(Q^2)$ gives $\overline{G}'(q^2) = (1 + Q^2/m_\pi^2) \overline{G}(Q^2)$. Since, Eq. (8) gives $\overline{G}(q^2)$ to lowest chiral order (*i.e.* up to corrections of order m_π^2/Λ^2 or Q^2/Λ^2 but including Q^2/m_π^2 to all orders) we immediately have the lowest order expression for $\overline{G}'(Q^2)$:

$$\overline{G}'(Q^2) = g_A \left(1 + \frac{2Q^2}{3m_\pi^2}\right) + \mathcal{O}(Q^2/\Lambda^2, m_\pi^2/\Lambda^2). \quad (15)$$

From the form of Eq. (15) it is immediately clear that for any finite quark mass, $\overline{G}'(0) = \overline{G}(0) = g_A$. Thus, in principle one can use either A_μ^a or A_μ^a to extract g_A .

The form of Eq. (15) shows that the current associated with A_μ^a is *not* delocalized in the sense that a finite

fraction of the integrated strength is at distance scales of order $1/m_\pi$. Were it so delocalized, all positive moments of the distribution would diverge as $m_\pi \rightarrow 0$ with $\langle r^{2n} \rangle_{A'} \sim 1/m_\pi^{2n}$. These moments are given by

$$\langle r^{2n} \rangle_{A'} = \prod_{j=1,n} (2+4j) (-1)^n \frac{d^n \overline{G}'(Q^2)}{d(Q^2)^n}, \quad (16)$$

which from Eq. (15) implies that

$$\begin{aligned} \langle r^2 \rangle_{A'} &= -\frac{4}{m_\pi^2} (1 + m_\pi^2/\Lambda^2) \\ \langle r^{2n} \rangle_{A'} &< \mathcal{O}(m_\pi^{-2n}) \text{ for } n > 1. \end{aligned} \quad (17)$$

Thus none of the higher moments diverge with m_π in a fashion consistent with a delocalized distribution.

One might ask how the r^2 moment could diverge with $1/m_\pi^2$ if the distribution is localized. The answer is simple: the spatial distribution $\rho_{A'}(r)$ could be of the following form:

$$\begin{aligned} \rho_{A'}(r) &= g_A \Lambda^3 f_1(\Lambda r) + \frac{\Lambda^6}{m_\pi^4} \partial_r f_2(\Lambda^2 r/m_\pi) \\ \text{with } \int dr 4\pi r^2 f_1(r) &= 1 \end{aligned} \quad (18)$$

where $f_2(x)$ goes to zero as $x \rightarrow \infty$ faster than any power law. The first term in Eq. (18) integrates to the full value of g_A while the second term integrates to zero due to the derivative. Moreover, the second term, although of large overall scale, is of very short range. Thus it leads to the moments of Eq. (17). It is clear that as the chiral limit is approached the second term remains sizable only over very short distances and cannot be associated with delocalization.

On the other hand, it is straightforward to see that if one uses A_μ^a to calculate g_A on a finite volume one gets exactly the same result as if one uses A_μ^a . To see this consider

$$\begin{aligned} g_A^{\text{Vol}} &\equiv \langle \vec{p}', I_3', s' | \int_{\text{box}} d^3x A_3^3(\vec{x}, t) | \vec{p}, I_3, s \rangle \\ &= \langle \vec{p}', I_3', s' | \int_{\text{box}} d^3x \left(1 + \frac{\partial^\mu \partial_\mu}{m_\pi^2}\right) A_3^3(\vec{x}, t) | \vec{p}, I_3, s \rangle. \end{aligned} \quad (19)$$

The second equality shows that the matrix element of A_3^3 has two parts—the matrix element of A_3^3 and a total derivative term. This second term can be written as

$$\begin{aligned} &\langle \vec{p}', I_3', s' | \int_{\text{box}} d^3x \frac{\partial^2}{\partial t^2} A_3^3(\vec{x}, t=0) | \vec{p}, I_3, s \rangle \\ &= \frac{L^3}{m_\pi^2} \langle \vec{p}', I_3', s' | \int_{\text{surface}} d^2x \hat{n} \cdot \vec{\nabla} A_3^3(\vec{x}, t=0) | \vec{p}, I_3, s \rangle, \end{aligned} \quad (20)$$

where $\hat{n}$ is an outward oriented normal vector; the time derivative vanishes since our matrix element is evaluated for an energy eigenstate and the divergence theorem is used to replace the volume integral by a surface integral.

On the other hand, the standard boundary conditions implies that $A_3^3(\vec{x}, t) = A_3^3(\vec{x} + L\hat{x}, t) = A_3^3(\vec{x} + L\hat{y}) = A_3^3(\vec{x} + L\hat{z})$ which in turn means that $\nabla A_3^3(\vec{x}, t=0)$ on one wall of the box is identical to $\nabla A_3^3(\vec{x}, t=0)$ on the opposite wall. Since the outward normal is opposite on opposite walls the integral vanishes. Only the first term of Eq. (19) contributes and thus one has $g_A^{\text{Vol}} = g_A^{\text{Vol}}$.

This completes the demonstration. The current A'_μ unlike A_μ remains localized in the chiral limit and hence cannot suffer from large finite volume effects due to delocalization. However, it gives the same result for g_A on a finite system as the original current provided the standard boundary conditions are imposed. Thus g_A calculated on a finite volume from A'_μ also cannot have large finite volume effects due to delocalization. The proceeding demonstration shows that the intuition that the delocalization of the axial current implies large finite volume effects is not correct.

It is useful to understand in a more heuristic manner why this intuition fails. Let us recall that the underlying reason for concluding the current was delocalized in the chiral limit was the rapid variation of $G(Q^2)$ for small pion masses associated with the pion pole. To see what is happening, it is sensible to restate the intuition that large finite volume effects might be expected in terms of momentum space. If one visualizes restricting the nucleon to a box, one might think that the states will not have well-defined momenta and the characteristic momenta in the state will be of order $1/L$. In such a case the matrix element would have components of momentum with $Q \sim 1/L$ and the rapid variation of $G(Q^2)$ might be expected to yield large finite volume effects unless $m_\pi L \gg 1$. However, this intuition is faulty. In particular, the perception that the momentum is not well-defined, while valid for boundary conditions in which the fields vanish on the walls, is *not* valid with the standard boundary conditions (periodic for bosons; antiperiodic for fermions). As discussed above, however, the standard boundary conditions are compatible with states of good momentum. The effect of imposing the standard boundary conditions does not mix momenta. Instead these boundary conditions allow momentum eigenstates but restrict the allowable momenta. Thus, with these boundary conditions one can have $\vec{Q} = 0$ exactly, and the fact that matrix elements vary rapidly with $\vec{Q}$ near $\vec{Q} = 0$ becomes irrelevant. The heuristic argument why delocalization in the chiral limit does not lead to large finite volume effects depends critically on the use of the standard boundary conditions. This is consistent with the more formal demonstration above in which standard boundary conditions were also essential in showing that the surface integral vanishes.

It is also useful to understand the failure of the intuition that for finite volumes and a delocalized source one expects to lose some of the strength out of the box and hence to have large finite volume effects. Again the

boundary conditions provide the explanation. The key point is that with these boundary conditions, a nucleon in the box is equivalent to an infinite system with a cubic array (with a lattice spacing of L) of boxes each containing a single nucleon. Thus, while strength can be lost from a box into its neighbors due to delocalization, a compensating strength may be gained from strength escaping into the box from its neighbors.

In summary, it has been shown that the delocalization of the axial current discussed in Ref. [3] does not lead to large finite volume effects. This was shown explicitly by constructing a current which is manifestly not delocalized and which leads to the same finite volume value for g_A as the standard current. The demonstration depended on the imposition of the standard boundary conditions. It was argued at a heuristic level why it is not surprising that delocalization does not yield large finite volume effects provided these boundary conditions are imposed.

This result leaves open the question of why present lattice calculations substantially underpredict g_A . Recent lattice results do suggest large finite volume effects [2]. This does not contradict the present result. Here it is only shown that the delocalization effect will not lead to large finite volume corrections. Of course, other effects can cause big finite volume effects. Given the relatively small volumes used for the present generation of lattice calculations it is not too surprising that large finite volume effects are present. However, there is also no reason to connect them with the delocalization effect of Ref. [3]. Ultimately, reliable lattice calculations should become available on larger lattices and the payoff of the chiral and infinite volume limits can be explored in the calculation of g_A . From the arguments presented here, one expects such calculations to indicate that the combined limit is uniform.

TDC gratefully acknowledges discussions with J-W Chen, X. Ji and R. L Jaffe. This work was supported in part by the U.S. Department of Energy.

-
- [1] M. Fukugita, Y. Kuramashi, M. Okawa and A. Ukawa, Phys. Rev. Lett. **75**, 2092 (1995); S.J. Dong, J.-F. Lagaë and K.F. Liu, Phys. Rev. Lett. **75**, 2096 (1995); K.F. Liu, S.J. Dong, T. Draper and J.M. Wu, Phys. Rev. **D49**, 4755 (1994); M. Göckeler et al, Phys. Rev. **D53**, 2317 (1996); S. Capitani et al, in Proc. Lattice 98, Nucl. Phys. B (Proc. Suppl.) **73**, 294 (1999); S. Capitani et al, Proc. DIS 99, Nucl. Phys. B (Proc. Suppl.) **79**, 548 (1999); S. Güsken et al, Phys. Rev. **D59**, 114502 (1999); T. Blum, S. Ohta, S. Sasaki, *Domain Wall Fermion Calculation of Nucleon g_A/g_V* , contributed to the 18th International Symposium on Lattice Field Theory (Lattice 2000). Published in Nucl. Phys. B (Proc. Suppl.) **94**, 295 (2001) [hep-

- lat/0011011]; R. Horsley (UKQCD and QCDSF Collaborations), *Determining g_A Nonperturbatively $\mathcal{O}(A)$ with Improved Wilson Fermions*, Talk presented at the 18th International Symposium on Lattice Field Theory (Lattice 2000). Published in Nucl. Phys. B (Proc. Suppl.) **94**, 307 (2001) [hep-lat/0010059].
- [2] Shoichi Sasaki, Tom Blum, Shigemi Ohta and Kostas Orginos, *Nucleon axial charge from quenched lattice QCD with domain wall fermions and improved gauge action*, hep-lat/0110053.
 - [3] R. L. Jaffe, *Delocalization of the axial charge in the chiral limit*, hep-ph/0108015.
 - [4] This issue was raised briefly in the classic paper on the phenomenology of the Skyrme model by Adkins, Nappi and Witten (G. S. Adkins, C. R. Nappi and E. Witten, Nucl. Phys. **B228**, 552 (1983)) in which this effect is responsible for a factor of 3/2 in the expression for the g_A in the chiral limit. More explicit discussions can be found in the context of chiral bag models in R. L. Jaffe, “The Bag”, in *Pointlike Structures Inside and Outside Hadrons, Proceedings of the 1979 Erice School in Subnuclear Physics*, A. Zichichi, ed. (Plenum, New York, 1981) and in the context of quark-based hedgehog soliton in appendix B of T. D. Cohen and W. Broniowski, Phys. Rev. **D34**, 3472 (1986).
 - [5] See, for example, T. P. Cheng and L. F. Lee, *Gauge Theory of Elementary Particle Physics* (Oxford, London, 1984), p. 153.
 - [6] M. Goldberger and S. B. Treiman, Phys. Rev. **110** 1178 (1958).